

TO OPEN...
CUT HERE!

RESEALABLE
ZIPPER
BAG

Hi-Yield®

10% CARBARYL Garden & Pet Dust



*Controls Listed Insects On Listed Vegetables, Berries,
Grapes, Flowers, Shrubs And Lawns.*

- Ready To Use 10% Dust.
- Kills Listed Insects On Fruits, Vegetables, Ornamentals, Lawns, Trees And Shrubs.
- For Use On Dogs And Cats.

NET WEIGHT 4 LBS. (1.8 KG)

**FOR HOMEOWNER
USE ONLY.**

ACTIVE INGREDIENT:
Carbaryl(1-naphthyl) N-methylcarbamate..... 10.00%
INERT INGREDIENTS:..... 90.00%
TOTAL:..... 100.00%

**KEEP OUT OF REACH
OF CHILDREN**

CAUTION

See Back Panel For First Aid,
Precautionary Statements and
Directions For Use.

31125

PRECAUTIONARY STATEMENTS HAZARDS TO HUMAN AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or if absorbed through the skin. Avoid contact with skin, eyes or clothing. Causes moderate eye irritation. Harmful if inhaled. Avoid breathing dust. Remove contaminated clothing and wash clothing before reuse. Wash thoroughly with soap and water after handling. Keep out of reach of children and domestic animals. Avoid contamination of feed and feed stuffs. Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

Food utensils such as teaspoons and tablespoons should not be used for food purposes after use with pesticides.

USER SAFETY REQUIREMENTS

- Wear long sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when mixing and applying this product.
- Change clothing as soon as possible after use.
- Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.
- Do not allow children or pets to contact treated area until dust has settled.

FIRST AID N-methyl Carbamate

If swallowed:	<ul style="list-style-type: none">• Call a Poison Control or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by the Poison Control Center or doctor.• Do not give anything by mouth to an unconscious person.
If inhaled:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.• Call a Poison Control Center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a Poison Control Center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a Poison Control Center or doctor for treatment advice.

Have the product container or label with you when calling a Poison Control Center or doctor, or going for treatment. You may also contact 1-800-222-1222 or 1-800-900-4044 for emergency medical information.

NOTE TO PHYSICIAN AND VETERINARIANS: Carbaryl is an N-methyl carbamate that inhibits cholinesterase. Atropine is antidotal. DO NOT use 2-PAM, opiates or cholinesterase-inhibiting drugs.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic invertebrates. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment and disposal of equipment washwaters. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

NOTE: Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith. Voluntary Purchasing Groups, Inc. warrants this product to conform to the chemical description on the label and for the purposes stated on the label. The limit of any liability incurred shall be the purchase price paid by the User or Buyer.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Wearing household latex or rubber gloves, use a dust applicator such as a shaker can, dust gun, puffer or other bulbous duster, rotary duster or other appropriate duster as available. Dust lightly to cover upper and lower leaf surfaces with a thin, even film or dust. Apply when Insects or their damage first appear. Apply when air is calm to avoid drift and contact with eyes and skin. Start applying at the farthest corner or the treatment area and work backward to avoid contact with dusted surfaces. Allow dust to settle in treated areas before reentering. Repeat at 7 to 10 day intervals as necessary for proper Insect control. For Plants above chest high, consider using a liquid Carbaryl product.

For residential use on Ornamentals, Flowers, Shrubs, Lawns, Home Garden, Vegetables and Pets.

ORNAMENTALS, FLOWERS, SHRUBS Roses, Azaleas, Camellias, Gardenias, Chrysanthemums, Rhododendron, Arborvitae, Boxwoods, Junipers, Holly Carnations, Gladiola, Lilac and Evergreens

To kill *Elm Leaf, Rose and Apple Aphids, Rose Slugs, Birch Leaf, Boxwood and Oak Leaf Miners, Mealy Bugs, Leaf Rollers, Plant Lace and Boxelder Bugs, Bagworms, Elm Leaf, Blister, Japanese, June and Willow (Crawlers), Orange Tortrix, Mimosa Webworms and Gypsy Moths:* Dust lightly to cover upper and lower leaf surfaces with a thin, even film of dust. Apply when Insects or their damage first appear. Repeat at 7 to 10 day intervals as necessary for proper Insect control. Do not use on Boston Ivy since injury may occur.

LAWNS

To kill *Chinch Bugs, Sod Webworms (Lawn Moths), Fall Armyworms, Cutworms, Ants, Fleas, Mosquitoes, Earwigs, Leafhoppers and Millipedes:* Apply 2 pounds per 1,000 square feet. For best results, mow lawn and apply immediately after rain or watering, and do not water for at least 2 days after application. Repeat at 2-3 week intervals if necessary. In Florida use 1 lb. per 400 sq. ft. for Chinch Bugs.

IMPORTED FIRE ANTS

For use as a mound treatment to control Imported Fire Ants, apply 1 to 2 ounces of product over and around each Fire Ant mound. Do not irrigate following application. Lightly disturb mound and immediately treat exposed mound surface. For best results apply early in the morning or late evening hours when Ants are least active. Treat new mounds as they appear.

VEGETABLES

The Pre-Harvest Interval (PHI) indicates the number of days between the last application and Harvest. The PHI is listed in the () immediately after each crop heading. Do not repeat applications more often than every 7 days.

BEANS (3)

To kill *Mexican Bean, Bean Leaf, Cucumber, Flea and Japanese Beetles, Leafhoppers, Velvet Bean Caterpillar, Western Bean Cutworm, Armyworms, Cutworms, Stink Bugs and Tarnished Plant Bugs* on Green Beans, Lima Beans, Snap Beans, Cowpeas and Black-eyed Peas. Use 0.25 lbs. per 1,000 square feet. When Mexican Bean Beetle is the only problem, a rate of 0.12 per 1,000 square feet is sufficient.

CARROTS, HEAD LETTUCE, RADISHES, PARSNIPS, AND TURNIPS (Roots) (7)

To kill *Armyworms, Cabbage Worms, Corn Earworms, Tarnished Plant Bugs and Stink Bugs* on Lettuceheads for Corn Earworms. Use 0.12 to 0.5 lbs. per 1,000 square feet.

ENDIVE, KALE, LEAF LETTUCE, MUSTARD GREENS, SPINACH AND TURNIPS (Tops) (14-15)

To kill *Armyworms, Cabbage Worms, Corn Earworms, Tarnished Plant Bugs and Stink Bugs.* Use 0.12 to 0.5 lbs. per 1,000 square feet.

CUCUMBER, MELONS AND SQUASH (3)

To kill *Pickleworms, Melon Worms, Cucumber and Flea Beetles, Leafhoppers and Squash Bugs.* Avoid excessive applications. Do not use on Watermelons in Florida. Use 0.25 lbs. per 1,000 square feet.

GARDEN PEAS (3)

To kill *Leafhoppers, Colorado Potato Beetles, Armyworms and Grasshoppers.* Use 0.12 to 0.25 lbs. per 1,000 square feet.

POTATO (7) TOMATO, EGGPLANT AND PEPPERS (3)

To kill *Colorado Potato Beetles, Flea Beetles, Leafhoppers, European Corn Borers, Fall Armyworms, Lace Bugs, Tomato Fruitworms and Hornworms, Tarnished Plant Bugs and Stink Bugs.* Use 0.12 lbs. per 1,000 square feet for Colorado Potato Beetles. All other insects use 0.25 to 0.5 lbs. per 1,000 square feet.

OKRA (3)

To kill *Corn Earworms and Stink Bugs.* Use 0.12 to 0.25 lbs. per 1,000 square feet.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER AND KOHLRABI (3)

To kill *Flea Beetles, Harlequin Bugs, Armyworms, Cabbage Worms and Corn Earworms.* To avoid possible injury, do not apply when foliage is wet or when rain or excessive humidity is expected during the next 2 days. Use 0.12 to 0.25 lbs. per 1,000 square feet.

GRAPES (7)

To kill *Grape Leaf Folder, Grape Leafhopper and Redbanded Leaf Roller.* Use 0.25 to 0.5 lbs. per 1,000 square feet.

STRAWBERRIES (7)

To kill *Meadow Spittle Bug, Strawberry Leaf Roller and Strawberry Weevil.* Use 0.25 to 0.5 lbs. per 1,000 square feet. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other Carbaryl labels.

DOGS AND CATS

Use only on Dogs and Cats. Consult a veterinarian before using this product on debilitated, medicated, aged or nursing animals. To control Fleas, Ticks and Lice. Use product only outside or in a well ventilated area. Wearing household latex or rubber gloves, lightly dust entire animal avoiding pet's eyes, nose, mouth and genital areas. Apply also to the feet between toes and on legs. Rub or brush pet's hair to work dust down to the skin. Do not use this or any pesticide on debilitated, medicated, aged or nursing animals. Do not use on Kittens or Puppies under 12 weeks of age. Do not repeat application for 30 days. Lightly dust bedding and doghouse using a shaker can or other dust applicator. Dust in and around sleeping quarters and all areas where Ticks and Fleas occur. DO NOT USE IN HOMES. DO NOT APPLY TO RUGS AND CARPETS AS STAINING MAY OCCUR. DO NOT REAPPLY WITHIN 30 DAYS OF EACH TREATMENT.

Sensitivities may occur after using ANY pesticide products for pets. If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

Do not use this product on animals simultaneously or within 30 days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides, or chemicals. However, flea and tick collars may be immediately replaced.

STORAGE AND DISPOSAL

STORAGE: Store in original container and place in a locked storage area.

DISPOSAL: If Empty: Do not reuse this container. Place in trash or offer for recycling if available. **If Partly Filled:** Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

VOLUNTARY PURCHASING GROUPS, INC.

P. O. BOX 460 • HWY. 56 WEST

BONHAM, TEXAS 75418

Visit Us At: www.v-p-g.com

EPA Reg. No. 7401-166

EPA Est. No. 7401-TX-1



4-07